

Abstract

Dark matter (DM) is known to interact gravitationally. If it is ultralight then there will be unique macroscopic effects that may be probable in the same environments that we see the evidence for DM. I will discuss how ultralight DM can be fermionic, evading the Tremaine-Gunn bound, and the new relevant constraints including those from supermassive black holes. Finally, I will present a specific model that addresses some interesting hints/anomalies in terms of early supermassive black hole formation, ultrafaint dwarf galaxies, and possible gravitational wave signatures.

Connecting the Extremes: A Story of Supermassive Black Holes and Ultralight Dark Matter



Peter B. Denton



MIT

April 11, 2022



[1904.09242](#) (PRL) w/ Hooman Davoudiasl

[2008.06505](#) (PRD) w/ Hooman Davoudiasl and David McGady

[2109.01678](#) (PRL) w/ Hooman Davoudiasl and Julia Gehrlein



Brookhaven™
National Laboratory

Speaking from occupied [Massachusetts](#) land

Outline

Superradiance, M87*, ultralight bosons, fuzzy DM

1. Superradiance probes the existence of ultralight bosons
2. M87* provides constraints
3. Relevant for fuzzy DM

Ultralight fermionic DM

1. Fermionic dark matter **can** be lighter than 100 eV
2. New limits arise from LHC, cosmic rays, black holes, ...
3. Strong gravity becomes important
4. How many species of particles are there?

A model at 10 keV

1. Early production of supermassive black holes
2. Axion dark matter
3. Gravitational waves




Superradiance, M87*, Fuzzy DM



Event Horizon Telescope: [ApJL 875 L1 \(2019\)](#)


What is this good for?

Black holes seem to follow $r \propto M$ over a huge range of masses



What is this good for?

Black holes seem to follow $r \propto M$ over a huge range of masses



Superradiance

Rotating BHs will create particles on-shell out of the vacuum:
Extracts angular momentum

Y. Zeldovich JETP Lett. 14, 180 (1971)

Conceptually similar to Hawking and Unruh radiation

Phenomenologically:

BHs can constrain the *existence* of bosons, independent of coupling

A. Arvanitaki, et al. [0905.4720](#)

A cloud of particles forms around the BH \Rightarrow no fermions¹

Care is needed for axions²

¹See slide [16](#)

²See slide [50](#)

Superradiance

Boson cloud growth rate:

$$\Gamma_0 = \frac{1}{24} a^* G^8 M^8 \mu_B^9, \quad \Gamma_1 = 4 a^* G^8 M^8 \mu_B^7$$

Leading to an occupation number after spinning down Δa^* :
 $a^* \equiv J/GM^2 \in [-1, 1]$

$$N = GM\Delta a^*$$

Superradiance depletes the spin of a BH if:

$$e^{\Gamma_B \tau_{\text{BH}}} > N$$

$\tau_{\text{BH}} \sim$ time to spin the BH back up

Wavelength has to enter into the ergosphere:

$$\mu_B > \Omega_H$$


Angular velocity:

$$\Omega_H \equiv \frac{1}{2GM} \frac{a^*}{1 + \sqrt{1 - a^{*2}}}$$

Only include dominant $m = 1$ spherical harmonic mode

M. Baryakhtar, R. Lasenby, M. Teo [1704.05081](#)

Spin



EHT: [ApJL 875 L5 \(2019\)](#)

- ▶ EHT can infer the spin
- ▶ Some degeneracies with disk properties
- ▶ EHT (conservative): $|a^*| \gtrsim 0.5$
- ▶ Circularity: No real power yet

C. Bambi, et al. [1904.12983](#)

- ▶ Twisted light: $|a^*| = 0.9 \pm 0.05$ at 95%

F. Tamburini, B. Thidé, M. Valle [1904.07923](#)

rules out $a^* = 0$ at 6σ

If a BH with large $|a^*|$ is measured, it could not have spun down much



Time scale

Astrophysics can spin the BH back up, possibly faster than superradiance

- ▶ From the Eddington limit, $\tau_{\text{Salpeter}} \sim 4.5 \times 10^7$ yrs
- ▶ EHT: $\dot{M}_{\text{M87}^*} / \dot{M}_{\text{Edd}} \sim 2 \times 10^{-5}$
- ▶ Mergers: one $\sim 10^9$ yrs ago with a much smaller galaxy
A. Longobardi, et al. [1504.04369](#)
- ▶ μ_B constraint has very weak dependence: $\tau_{\text{BH}}^{-1/7}$ or $\tau_{\text{BH}}^{-1/9}$

We take $\tau_{\text{BH}} = 10^9$ yrs


Past ultra light boson constraints



Spin-1 constraints

M. Baryakhtar, R. Lasenby, M. Teo [1704.05081](#)


New constraints from M87*



Bosons with masses in the regions in color are ruled out.

Superradiance Spin-down



Different spherical harmonic modes leads to different maximum spins



Vector (scalar) in bold (dotted) for $\mu_B = 10^{-12}$ eV

M. Baryakhtar, R. Lasenby, M. Teo [1704.05081](#)

How to detect ultra light bosons with superradiance



Vector with $\mu_B = 10^{-12}$ eV

$\sigma_{a^*} \sim 0.3, \sigma_M/M \sim 10\%$

M. Baryakhtar, R. Lasenby, M. Teo [1704.05081](#)

Superradiance conclusions

- ▶ Superradiance is a powerful probe of ultralight bosons
- ▶ Constraints from M87* relevant for fuzzy DM
- ▶ Discovering ultralight bosons is hard

Ultralight Fermionic DM

Dark matter: what we know

Astrophysically/gravitationally: lots

Particle nature:

- ▶ Coupling to SM/self? Could be zero (other than gravity)
- ▶ Heavier than $\sim 100 M_{\odot}$ leads to tidal disruption effects
- ▶ Lighter than $\sim 10^{-22}$ eV, at $v \sim 10^{-3}$, Compton wavelength is too big
 - ▶ Core/cusp suggests $\sim 10^{-22} - 10^{-21}$ eV
- ▶ Fermionic DM lighter than ~ 100 eV can't be squeezed into a galaxy

S. Tremaine, J. Gunn [PRL 42, 407 \(1979\)](#)

Light fermionic dark matter

Light fermionic dark matter $m < 100$ eV can't be squeezed into galaxies

Two issues:

1. Getting light thermal population into low momentum states is difficult
2. Pauli exclusion principle

S. Tremaine, J. Gunn [PRL 42, 407 \(1979\)](#)

Focus on #2

Light fermionic dark matter

Light fermionic dark matter $m < 100$ eV can't be squeezed into galaxies

Two issues:

1. Getting light thermal population into low momentum states is difficult
2. Pauli exclusion principle

S. Tremaine, J. Gunn [PRL 42, 407 \(1979\)](#)

Focus on #2

Modern treatments find that the limit is

▶ 100 eV

C. Di Paolo, et al. [1704.06644](#)

▶ 190 eV (2σ)

D. Savchenko, A. Rudakovskiy [1903.01862](#)

▶ 130 eV (2σ)

J. Alvey, et al. [2010.03572](#)

Evading Tremaine-Gunn

Dark matter could be composed of many different species

The correct bound on light fermionic DM:

$$N_F \gtrsim \left(\frac{100 \text{ eV}}{m} \right)^4$$

- ▶ One power: lighter DM requires more species
- ▶ Three powers: phase space

So 1 eV fermionic DM is possible if there are $N_F \gtrsim 10^8$ species.

“Model”

Different species can be degenerate:

$$\mathcal{L} \supset -m \sum_{i=1}^{N_F} \bar{\chi}_i \chi_i$$

Perhaps $SU(\sqrt{N_F})$ which leads to quasi-degenerate states:

$$\frac{m_i - m_j}{m_1} \sim \frac{\lambda^2}{16\pi^2} \log \frac{m_1}{\Lambda}$$

m_1 is the lightest mass

L. Randall, J. Scholtz, J. Unwin [1611.04590](#)

Perhaps Kaluza-Klein modes: Constraint is more complicated

Extrapolation!

Let's extrapolate this as far as possible!


$$m \gtrsim 10^{-22} \text{ eV} \Rightarrow N_F \gtrsim 10^{96}$$

How many DM particles would there be in a galaxy in this case?

Dwarf spheroidals have $\sim 10^{96}$ DM particles if $m \sim 10^{-22}$ eV

Below this the fourth power scaling law drops to $N_F \gtrsim \left(\frac{100 \text{ eV}}{m}\right)^4$

No more Pauli exclusion



Extrapolation!

Let's extrapolate this as far as possible!


$$m \gtrsim 10^{-22} \text{ eV} \Rightarrow N_F \gtrsim 10^{96}$$

How many DM particles would there be in a galaxy in this case?

Dwarf spheroidals have $\sim 10^{96}$ DM particles if $m \sim 10^{-22}$ eV

Below this the fourth power scaling law drops to $N_F \gtrsim \left(\frac{100 \text{ eV}}{m}\right)^4$

No more Pauli exclusion



Too many species

Claim:

10^{96} species is Too Many

SM has 10^2 species

From now it doesn't matter:

1. if the species are DM,
2. if they are fermions,
3. if their masses are degenerate

Too many species

Claim:

10^{96} species is Too Many

SM has 10^2 species

From now it doesn't matter:

1. if the species are DM,
2. if they are fermions,
3. if their masses are degenerate

Gravitational effects are suppressed by M_P , but enhanced by N

$$\sum_i^N \sigma_i \sim N \frac{E^2}{M_P^4}$$

Cosmic ray constraints

Highest energy collisions recorded are UHECRs

Telescope Array and the Pierre Auger Observatory see a suppression at $10^{19.5}$ eV

O. Deligny for TA and Auger [2001.08811](#)


Could be photo-pion production (GZK)

K. Greisen [PRL 16, 748 \(1966\)](#)

G. Zatsepin, V. Kuzmin [JETP Lett. 4, 78 \(1966\)](#)

Could be end of sources

See e.g. R.A. Batista, et al. [1903.06714](#)




Cosmic ray constraints

Can use cosmic rays to constrain large number of species

1. As N increases, $BR(pp \rightarrow \chi\chi) \rightarrow 1$
2. Showers would be reconstructed at a lower energy
3. There would appear to be a suppression to the flux
4. No suppression is seen below $E_{\text{LAB}} \sim 10^{19.5}$ eV ($\sqrt{s} = 250$ TeV)


$$N \lesssim 4 \times 10^{68} \quad \text{for} \quad m \lesssim 100 \text{ TeV}$$



Lower energy, better precision

- ▶ Searches for monojets
 - ▶ Detected 245 events with $E_T^{miss} > 1$ TeV
 - ▶ Expected 238 ± 23
 - ▶ Mostly $Z \rightarrow \nu\nu$ with ISR or brem
- ATLAS 1711.03301
- ▶ $G \rightarrow \chi\chi$ looks the same
 - ▶ Include 3-body $(4\pi)^{-3}$ factor

$$N \lesssim 10^{62} \quad \text{for} \quad m \lesssim 500 \text{ GeV}$$




100 TeV will improve by $\sim 2+$ orders of magnitude

BH evaporation

- ▶ $t_{evap} \sim \frac{10^{67}}{N} \left(\frac{M_{BH}}{M_{\odot}} \right)^3 \text{ yr}$
- ▶ We assume that $M_{BH} \sim 10M_{\odot}$ have been around for $\sim 10^9 \text{ yr}$
- ▶ $10M_{\odot} \rightarrow T_{BH} \sim 10^{-11} \text{ eV}$

$$N \lesssim 10^{61} \quad \text{for} \quad m \lesssim 10^{-11} \text{ eV}$$



Fermionic DM can be as light as $\sim 10^{-13}$ eV


Need $\sim 10^{61}$ quasi-degenerate species

These constraints apply regardless of whether it is
DM, fermionic, or quasi-degenerate

Low energies but high densities

- ▶ New states populated via gravity in the early universe
- ▶ Don't want $\rho_\chi \gtrsim \rho_\gamma$
- ▶ $\rho_\chi/\rho_\gamma \sim NT^3/M_P^3$
- ▶ Implies a maximum reheat temperature
- ▶ BBN requires $T_{rh} \gtrsim 10$ MeV

$$N \lesssim 10^{63} \quad \text{for} \quad m \lesssim 10 \text{ MeV}$$




Supernovae

Low energies but high densities and more measurements

- ▶ Neutrino production $\sigma_\nu \sim E^2 G_F^2$
- ▶ Dark sector production $\sigma_\chi \sim N E^2 / M_P^4$
- ▶ Can't have a significant amount of energy to dark sector
- ▶ $N \lesssim G_F^2 M_P^4$

$$N \lesssim 10^{66} \quad \text{for} \quad m \lesssim 100 \text{ MeV}$$




Superradiance with bosons

Narrow applicability range, apply down to $N_B = 1$ for bosons

- ▶ Power law for small masses m^{-9}
- ▶ Exponential for large masses
- ▶ Conservatively take constraints on $S = 0$
- ▶ Different regions are distinct constraints


H. Davoudiasl, [PBD 1904.09242](#)

M. Baryakhtar, R. Lasenby, M. Teo [1704.05081](#)



Superradiance with fermions

- ▶ Power law for small masses m^{-6}
- ▶ Exponential for large masses
- ▶ Conservatively take constraints on $S = \frac{1}{2}$
- ▶ Different regions are distinct constraints
- ▶ If $N_F \lesssim$ cloud occupation number, superradiance stops
 - ▶ Occupation number $\sim 10^{77}$ for stellar mass BH



Neutrino oscillations

If neutrinos get mass via usual seesaw, can write down:

$$\xi_i H^* \bar{\ell} \chi_i$$

leads to oscillations

$$P(\nu_\ell \rightarrow \chi_i) \sim \frac{\xi_i^2 \langle H \rangle^2}{m_\nu^2} \sin^2 \left(\frac{m_\nu^2 L}{4E} \right)$$

Assume $m_{\nu, \text{lightest}}$ is not too light

$$\langle H \rangle^2 / m_\nu^2 \sim 10^{24}$$

Neutrino oscillations

If neutrinos get mass via usual seesaw, can write down:

$$\xi_i H^* \bar{\ell} \chi_i$$

leads to oscillations

$$P(\nu_\ell \rightarrow \chi_i) \sim \frac{\xi_i^2 \langle H \rangle^2}{m_\nu^2} \sin^2 \left(\frac{m_\nu^2 L}{4E} \right)$$

Assume $m_{\nu, \text{lightest}}$ is not too light

$$\langle H \rangle^2 / m_\nu^2 \sim 10^{24}$$

$$P(\nu_\ell \rightarrow \chi) \sim N_F P(\nu_\ell \rightarrow \chi_i) \lesssim 0.1$$

$$N_F \xi_i^2 \lesssim 10^{-25}$$

To be competitive with LHC, need $\xi_i \gtrsim e^{-97}$
Instanton effects should suppress by $\sim e^{-100}$

L. Abbott, M. Wise [NPB 325, 687 \(1989\)](#)

R. Kallosh, et al. [hep-th/9502069](#)

P. Svrcek, E. Witten [hep-th/0605206](#)

H. Davoudiasl [2003.04908](#)

L. Hui, et al. [1610.08297](#)

Nucleon decay

One can write down this operator

$$\mathcal{O} \sim \frac{udd\chi_i}{M_P^2}$$

$$\Gamma(p \rightarrow \pi^+ + \chi) \sim N_F \frac{m_p^5}{M_P^4}$$

$$N_F \lesssim 10^{12} \quad \text{for} \quad m \lesssim 100 \text{ MeV}$$

If there is an associated global $U(1)$ charge, an instanton would suppress this rate by $e^{-200} \sim 10^{87}$

Strong gravity

Literature suggests that at $N \sim 10^{32}$ something happens with strong gravity at $m \sim 1$ TeV

G. Dvali [0806.3801](#)

I. Antoniadis, et al. [hep-ph/9804398](#)

S. Adler [PRL 44, 1567 \(1980\)](#)

N. Arkani-Hamed, S. Dimopoulos, G. Dvali [hep-ph/9807344](#)

X. Calmet, S. Hsu, D. Reeb [0803.1836](#)

G. Dvali, M. Redi [0905.1709](#)

A. del Rio, R. Durrer, S. Patil [1808.09282](#)

$N \sim 10^{32}$ species with $m \lesssim 1$ TeV may pull M_P to electroweak

According to Dvali or Adler:

$$G^{-1}(\mu) \sim G^{-1}(0) - Nm^2 \log \frac{\mu^2}{m^2}$$

$$G^{-1}(0) = M_P^2$$

This leads to

$$m\sqrt{N} \lesssim M_P$$

Calmet:


$$G^{-1}(\mu) \sim G^{-1}(0) - \frac{N\mu^2}{12\pi}$$

Fermionic dark matter conclusions

- ▶ The “number of species” axis for DM is interesting
- ▶ Fermionic DM can be as light as 10^{-13} eV with key constraints from BH lifetimes and the LHC
- ▶ Many similar constraints on the number of species from cosmic rays, LHC, BH lifetimes, BBN, and SNe
- ▶ More work to be done on this topic in many directions: pheno and theory

A model at 10 keV

Early supermassive black holes



K. Inayoshi, E. Visbal, Z. Haiman [1911.05791](#)

- ▶ Appears SMBHs are forming larger/faster than expected
- ▶ Some are known to be sub-Eddington
- ▶ Masses are probably underestimated
- ▶ Seems like they can't form conventionally

Ultralight dark matter hints

- ▶ Core-cusp might indicate fuzzy DM
 $\sim 10^{-21}$ eV

W. Hu, R. Barkana, A. Gruzinov [astro-ph/0003365](#)

D. Marsh [1510.07633](#)

L. Hui, et al [1610.08297](#)


- ▶ Ultra faint dwarf galaxies great place to look
- ▶ Baryonic feedback expected to be negligible

A. Lazar et al [2004.10817](#)

- ▶ Ultrafaint dwarf galaxies might prefer ultralight DM
- ▶ Strongest: Segue 1 prefers $\sim 10^{-19}$ eV
- ▶ See also Lyman- α

K. Rogers, H. Peiris [2007.12705](#)

[2109.01678](#)



K. Hayashi, E. Ferreira, H. Chan [2102.05300](#)

Simple start

- ▶ Evidence suggests stringy models have many $\mathcal{O}(100 - 1000)$ axions

P. Svrcek, E. Witten [hep-th/0605206](#)

A. Arvanitaki, et al [0905.4720](#)

- ▶ Expect decay constant similar to reduced Planck mass

$$f_a \sim \bar{M}_p \sim 2 \times 10^{18} \text{ GeV}$$

$$m_a \sim \frac{\mu_a^2}{f_a} \sim 10^{-20} \text{ eV} \left(\frac{\mu_a}{\text{keV}} \right)^2 \left(\frac{10^{17} \text{ GeV}}{f_a} \right)$$

Early black hole formation

Suppose the BH forms in the early universe somehow:

$$\frac{M_{\text{BH}}}{M_P^2} \sim R_{\text{BH}} \sim H^{-1}$$

$$H \sim \frac{T^2}{M_P}$$


$$M_{\text{BH}} \sim \frac{M_P^3}{T^2}$$

For $M_{\text{BH}} \sim 10^9 M_\odot$ we get $H \sim 10^{-19}$ eV and $T \sim 10$ keV

Matches with fuzzy DM set from near the Planck scale

Two sides of the same coin

Ultralight DM and early SMBH formation linked via first order phase transition



- ▶ $SU(3)_d$
- ▶ Heavy quarks $m_\Psi \sim f_a \sim 10^{17}$ GeV
- ▶ Confinement scale $\mu_a \sim 10$ keV
- ▶ This leads to a FOPT at $T \sim 10$ keV
- ▶ Drop of pressure allows SMBH formation
- ▶ PQ charges leads to dark quark-dark gluon couplings

$$\mathcal{L} \supset \frac{a}{f_a} G_{d\mu\nu} \tilde{G}_d^{\mu\nu}$$

P. Forcrand, M. D'Elia [1702.00330](#)

Under the rug

Dark gluons are populated in the early universe as glueballs
Contribute to N_{eff}

$$\Delta N_{\text{eff}} = \frac{4}{7} \left(\frac{11}{4} \right)^{4/3} \left(\frac{T_d}{T} \right)^4 N_{dG} \lesssim 0.3$$

Planck [1807.06209](#)

$$\Rightarrow T_d \lesssim 0.36T$$

Glueballs decay to new light states, redshift as radiation

$$\Rightarrow \Delta N_{\text{eff}} \sim 0.1$$

Other predictions: gravitational waves


FOPT produce GWs in various ways

1. vacuum bubble collisions
2. sound waves ✓
3. magnetohydrodynamic turbulence

We assume:

- ▶ Energy released in FOPT is $\alpha < 1$
- ▶ Velocity is large $\beta/H_* \sim 10^4$

A. Helmboldt, J. Kubo, S. Woude [1904.07891](#)




10 keV model conclusions

- ▶ Hints of anomalous early SMBH formation
- ▶ Hints of ultralight DM
- ▶ Dark SU(3) FOPT at 10 keV forms SMBHs
- ▶ With scale set by string theory, get axion DM
- ▶ Axion DM explains fuzzy DM hints
- ▶ Predict small correction to ΔN_{eff}
- ▶ Predict GWs for future PTAs

Thanks!

Backups

Spin dependence



Superradiance combinatorics

Assumed that generating N_F particles out of N_F species yields N_F distinct species

Just because a large number of particles spanning a large number of species are produced doesn't mean that they are actually different

The expected number of distinct species is

$$N_F \left[1 - \left(\frac{N_F - 1}{N_F} \right)^{N_F} \right] \rightarrow N_F \left(1 - \frac{1}{e} \right) \approx 0.63 N_F$$

Less than factor of two \Rightarrow we're good

Strong gravity: deviations

A running in G would lead to variations in gravity on different scales

$$\frac{\delta G}{G} \lesssim 10^{-9} \quad \text{for} \quad \ell \gtrsim 10^3 \text{ km} \rightarrow 10^{-13} \text{ eV}$$

P. Fayet [1712.00856](#)

S. Schlamminger, et al. [0712.0607](#)

This is not as strong as the 10^{32} arguments

Strong gravity: deviations

A running in G would lead to variations in gravity on different scales

$$\frac{\delta G}{G} \lesssim 10^{-9} \quad \text{for} \quad \ell \gtrsim 10^3 \text{ km} \rightarrow 10^{-13} \text{ eV}$$

P. Fayet [1712.00856](#)

S. Schlamminger, et al. [0712.0607](#)

This is not as strong as the 10^{32} arguments


At $N \sim 10^{60}$ and $m \sim 10^{-3}$ eV consistent with theory arguments on previous slide

$$\Rightarrow \frac{\delta G}{G} \sim 10^{-2} \quad \text{for} \quad \ell \sim 0.1 \text{ mm}$$

Close to current constraints

J. Lee, et al. [2002.11761](#)

Superradiance constraints with interactions



M. Baryakhtar, et al. [2011.11646](#)