

The best way to probe CP violation in the lepton sector is with long-baseline accelerator neutrino experiments in the appearance mode: the appearance of ν_e in predominantly ν_μ beams. Here we show that it is possible to discover CP violation with disappearance experiments only, by combining JUNO for electron neutrinos and DUNE or Hyper-Kamiokande for muon neutrinos. While the maximum sensitivity to discover CP is quite modest (1.6σ with 6 years of JUNO and 13 years of DUNE), some values of δ may be disfavored by $> 3\sigma$ depending on the true value of δ .

CP-Violation with Neutrino Disappearance

Peter B. Denton

Rice University

April 2, 2024

2309.03262



BrookhavenTM
National Laboratory



History


A little bit of my history:


1. Attended Rice, class of 2010
2. PhD Vanderbilt+FNAL, defended 2016
3. Postdoc Niels Bohr Institute
4. Scientist at Brookhaven, 2018-now


History


A little bit of my history:


1. Attended Rice, class of 2010
2. PhD Vanderbilt+FNAL, defended 2016
3. Postdoc Niels Bohr Institute
4. Scientist at Brookhaven, 2018-now
5. Andrew Long visited BNL in 2023
 - ▶ Discussed an early universe model to probe CPV in the neutrino sector
 - ▶ A colleague asked if there were other ways to probe CPV without new physics
 - ▶ I first said no, appearance measurements like $\nu_\mu \rightarrow \nu_e$ is the only way
 - ▶ Remembered there is another way

















Four known unknown in particle physics: all neutrinos

Atmospheric mass ordering

θ_{23} octant

Complex phase

Absolute mass scale

Four known unknown in particle physics: all neutrinos

Atmospheric mass ordering

θ_{23} octant

Complex phase

Absolute mass scale

Outline

1. Why CPV is interesting
2. Other non-standard probes of CPV
3. Relationship between appearance, disappearance, CP, T, CPT
4. Three ways to see why there is CPV information in disappearance
 - 4.1 Parameter counting
 - 4.2 Direct analytic calculation
 - 4.3 Numerical test
5. Role of the matter effect
6. Recommendation

Why is CPV interesting?

δ and CP violation

$$J_{CP} = s_{12}c_{12}s_{13}c_{13}^2s_{23}c_{23} \sin \delta$$

C. Jarlskog [PRL 55, 1039 \(1985\)](#)



δ and CP violation

$$J_{CP} = s_{12}c_{12}s_{13}c_{13}^2s_{23}c_{23} \sin \delta$$

C. Jarlskog [PRL 55, 1039 \(1985\)](#)



1. Strong interaction: no observed EDM \Rightarrow CP (nearly) **conserved**

$$\frac{\bar{\theta}}{2\pi} < 10^{-11}$$

J. Pendlebury, et al. [1509.04411](#)

2. Quark mass matrix: non-zero but **small** CP violation

$$\frac{|J_{CKM}|}{J_{\max}} = 3 \times 10^{-4}$$

CKMfitter [1501.05013](#)



3. Lepton mass matrix: ?

$$\frac{|J_{PMNS}|}{J_{\max}} < 0.34$$

PBD, J. Gehrlein, R. Pesles [2008.01110](#)

$$J_{\max} = \frac{1}{6\sqrt{3}} \approx 0.096$$

δ : what is it really?



δ : what is it not?

$\delta \not\Rightarrow$ Baryogenesis

The amount of leptogenesis is a function of:

1. the heavy mass scale
2. δ
3. α, β (Majorana phases)
4. CP phases in the RH neutrinos
5. ...

C. Hagedorn, et al. [1711.02866](#)


K. Moffat, et al. [1809.08251](#)

Measuring $\delta = 0, \pi$	$\not\Rightarrow$	no leptogenesis
Measuring $\delta \neq 0, \pi$	$\not\Rightarrow$	leptogenesis

δ, J : current status

Maximal CP violation is already ruled out:

1. $\theta_{12} \neq 45^\circ$ at $\sim 15\sigma$
2. $\theta_{13} \neq \tan^{-1} \frac{1}{\sqrt{2}} \approx 35^\circ$ at many (100) σ
3. $\theta_{23} = 45^\circ$ allowed at $\sim 1\sigma$
4. $|\sin \delta| = 1$ allowed



When δ and when J ?

If the goal is **CP violation** the Jarlskog invariant should be used
however

If the goal is **measuring the parameters** one must use δ

Given θ_{12} , θ_{13} , θ_{23} , and J , I can't determine the sign of $\cos \delta$ which is physical
e.g. $P(\nu_\mu \rightarrow \nu_\mu)$ depends on $\cos \delta$

Other non-standard CPV probes


1. Some information in solar due to loops in elastic scattering

V. Brdar, X-J. Xu [2306.03160](#)


2. Sub-GeV atmospherics

K. Kelly, et al. [1904.02751](#)

See also e.g. A. Suliga, J. Beacom [2306.11090](#)



Solar (no systematics): $\sim 0.5\sigma$



Atmospherics at DUNE: $< 2\sigma$

Appearance, disappearance, and CP

Appearance vs. Disappearance

Oscillation experiments can do
appearance or disappearance experiments:

Disappearance

K2K, MINOS, T2K, NO ν A
KamLAND, Daya Bay, RENO, Double CHOOZ
(Sort of) SNO, Borexino, SK-solar
JUNO, DUNE, HK

Appearance


T2K, NO ν A
OPERA
Atm ν_τ hints @ SK & IceCube
DUNE, HK

Neither appearance nor disappearance

SK-atm, IceCube



CP, T: Disappearance




Disappearance measurements are even eigenstates of CP

$$CP[P(\nu_e \rightarrow \nu_e)] = P(\bar{\nu}_e \rightarrow \bar{\nu}_e) \stackrel{CPT}{=} P(\nu_e \rightarrow \nu_e)$$

Assume that CPT is a good symmetry

CP, T: Appearance



Appearance measurements are not eigenstates of CP

Appearance and Disappearance, CP even and CP odd terms

Disappearance:

$$\begin{aligned} P(\nu_\alpha \rightarrow \nu_\alpha) &= 1 - 4|U_{\alpha 1}|^2|U_{\alpha 2}|^2 \sin^2 \Delta_{21} \\ &\quad - 4|U_{\alpha 1}|^2|U_{\alpha 3}|^2 \sin^2 \Delta_{31} \\ &\quad - 4|U_{\alpha 2}|^2|U_{\alpha 3}|^2 \sin^2 \Delta_{32} \\ &= P_{\alpha\alpha}^{CP+} \end{aligned}$$

$$\Delta_{ij} \equiv \Delta m_{ij}^2 L / 4E$$

Appearance and Disappearance, CP even and CP odd terms

Disappearance:

$$\begin{aligned} P(\nu_\alpha \rightarrow \nu_\alpha) &= 1 - 4|U_{\alpha 1}|^2|U_{\alpha 2}|^2 \sin^2 \Delta_{21} \\ &\quad - 4|U_{\alpha 1}|^2|U_{\alpha 3}|^2 \sin^2 \Delta_{31} \\ &\quad - 4|U_{\alpha 2}|^2|U_{\alpha 3}|^2 \sin^2 \Delta_{32} \\ &= P_{\alpha \alpha}^{CP+} \end{aligned}$$

Appearance:

$$\begin{aligned} P(\nu_\alpha \rightarrow \nu_\beta) &= -4\Re[U_{\alpha 1} U_{\beta 1}^* U_{\alpha 2}^* U_{\beta 2}] \sin^2 \Delta_{21} \\ &\quad - 4\Re[U_{\alpha 1} U_{\beta 1}^* U_{\alpha 3}^* U_{\beta 3}] \sin^2 \Delta_{31} \\ &\quad - 4\Re[U_{\alpha 3} U_{\beta 3}^* U_{\alpha 2}^* U_{\beta 2}] \sin^2 \Delta_{32} \\ &\quad \pm 8J_{CP} \sin \Delta_{21} \sin \Delta_{31} \sin \Delta_{32} \\ &= P_{\alpha \beta}^{CP+} + P_{\alpha \beta}^{CP-} \end{aligned}$$

$\Delta_{ij} \equiv \Delta m_{ij}^2 L / 4E$

Sign depends on α, β

Conventional Wisdom

1. Appearance is sensitive to CPV [True]

Conventional Wisdom

1. Appearance is sensitive to CPV [True]
2. Disappearance has no CPV sensitivity [False]

Conventional Wisdom

1. Appearance is sensitive to CPV [True]
2. Disappearance has no CPV sensitivity [False]
3. Any δ dependence in disappearance is in ν_μ not ν_e [Confusing/False]

$$\begin{pmatrix} c_{13}c_{12} & c_{13}s_{12} & s_{13}e^{-i\delta} \\ -c_{23}s_{12} - s_{23}s_{13}c_{12}e^{i\delta} & c_{23}c_{12} - s_{23}s_{13}s_{12}e^{i\delta} & s_{23}c_{13} \\ s_{23}s_{12} - c_{23}s_{13}c_{12}e^{i\delta} & -s_{23}c_{12} - c_{23}s_{13}s_{12}e^{i\delta} & c_{23}c_{13} \end{pmatrix}$$

Correct Statements

- ▶ Appearance is the best way to measure δ and CPV
 - ... given known oscillation parameters, systematics, and realistic experiments
 - ▶ Probes mostly $\sin \delta$ not $\cos \delta$
 - ▶ Don't need both ν and $\bar{\nu}$ (but systematics)
- ▶ Disappearance can measure δ
 - ▶ CPV can be discovered with only disappearance measurements
 - ▶ Probes mostly $\cos \delta$ not $\sin \delta$
 - ▶ Requires measurements of two flavors
 - ▶ "Works through unitarity" (as do nearly all oscillation measurements)

Parameter Counting

1. Four parameters in the PMNS matrix

Majorana phases are irrelevant

Parameter Counting

1. Four parameters in the PMNS matrix

Majorana phases are irrelevant

2. Disappearance experiments of one flavor can measure up to three amplitudes

- ▶ Electron neutrino row:

- ▶ KamLAND measured one
- ▶ Daya Bay/RENO measured a different one
- ▶ JUNO will measure all three

- ▶ Muon neutrino row:

- ▶ MINOS/T2K/NO ν A measured one

Parameter Counting

1. Four parameters in the PMNS matrix

Majorana phases are irrelevant

2. Disappearance experiments of one flavor can measure up to three amplitudes

- ▶ Electron neutrino row:

- ▶ KamLAND measured one
- ▶ Daya Bay/RENO measured a different one
- ▶ JUNO will measure all three

- ▶ Muon neutrino row:

- ▶ MINOS/T2K/NO ν A measured one

$$P(\nu_\alpha \rightarrow \nu_\alpha) = 1 - 4 \sum_{i>j} C_{ij}^\alpha \sin^2 \Delta_{ij}$$

$$C_{ij}^\alpha = |U_{\alpha i}|^2 |U_{\alpha j}|^2$$

$$|U_{\alpha i}| = \left(\frac{C_{ij}^\alpha C_{ik}^\alpha}{C_{jk}^\alpha} \right)^{1/4}$$

Parameter Counting

1. Four parameters in the PMNS matrix

Majorana phases are irrelevant


2. Disappearance experiments of one flavor can measure up to three amplitudes

- ▶ Electron neutrino row:

- ▶ KamLAND measured one
- ▶ Daya Bay/RENO measured a different one
- ▶ JUNO will measure all three

- ▶ Muon neutrino row:

- ▶ MINOS/T2K/NO ν A measured one



Daya Bay [1809.02261](#)

Parameter Counting

1. Four parameters in the PMNS matrix

Majorana phases are irrelevant


2. Disappearance experiments of one flavor can measure up to three amplitudes

- ▶ Electron neutrino row:

- ▶ KamLAND measured one
- ▶ Daya Bay/RENO measured a different one
- ▶ JUNO will measure all three


- ▶ Muon neutrino row:

- ▶ MINOS/T2K/NO ν A measured one



JUNO [1507.05613](#)

2309.03262



L. Zhan, et al. [0807.3203](#)

Rice University: April 2, 2024 20/29

Parameter Counting

1. Four parameters in the PMNS matrix
Majorana phases are irrelevant
2. Disappearance experiments of one flavor can measure up to three amplitudes
 - ▶ Electron neutrino row:
 - ▶ KamLAND measured one
 - ▶ Daya Bay/RENO measured a different one
 - ▶ JUNO will measure all three
 - ▶ Muon neutrino row:
 - ▶ MINOS/T2K/NO ν A measured one
3. Only two parameters for one flavor are independent

Parameter Counting

1. Four parameters in the PMNS matrix
Majorana phases are irrelevant
2. Disappearance experiments of one flavor can measure up to three amplitudes
 - ▶ Electron neutrino row:
 - ▶ KamLAND measured one
 - ▶ Daya Bay/RENO measured a different one
 - ▶ JUNO will measure all three
 - ▶ Muon neutrino row:
 - ▶ MINOS/T2K/NO ν A measured one
3. Only two parameters for one flavor are independent
4. Given good measurements of the ν_e and ν_μ disappearance, 4 independent parameters will be measured
 - ▶ Any row can be “simple” (e.g. $c_{12}c_{13}$, $s_{12}c_{13}$, ...) \Rightarrow no one row is ever enough
 - ▶ That is, CPV is physical and cannot depend on parameterization

Parameter Counting

1. Four parameters in the PMNS matrix
Majorana phases are irrelevant
2. Disappearance experiments of one flavor can measure up to three amplitudes
 - ▶ Electron neutrino row:
 - ▶ KamLAND measured one
 - ▶ Daya Bay/RENO measured a different one
 - ▶ JUNO will measure all three
 - ▶ Muon neutrino row:
 - ▶ MINOS/T2K/NO ν A measured one
3. Only two parameters for one flavor are independent
4. Given good measurements of the ν_e and ν_μ disappearance, 4 independent parameters will be measured
 - ▶ Any row can be “simple” (e.g. $c_{12}c_{13}$, $s_{12}c_{13}$, ...) \Rightarrow no one row is ever enough
 - ▶ That is, CPV is physical and cannot depend on parameterization
5. This is sufficient to constrain $\cos \delta$ and three mixing angles

Parameter Counting

1. Four parameters in the PMNS matrix
Majorana phases are irrelevant
2. Disappearance experiments of one flavor can measure up to three amplitudes
 - ▶ Electron neutrino row:
 - ▶ KamLAND measured one
 - ▶ Daya Bay/RENO measured a different one
 - ▶ JUNO will measure all three
 - ▶ Muon neutrino row:
 - ▶ MINOS/T2K/NO ν A measured one
3. Only two parameters for one flavor are independent
4. Given good measurements of the ν_e and ν_μ disappearance, 4 independent parameters will be measured
 - ▶ Any row can be “simple” (e.g. $c_{12}c_{13}$, $s_{12}c_{13}$, ...) \Rightarrow no one row is ever enough
 - ▶ That is, CPV is physical and cannot depend on parameterization
5. This is sufficient to constrain $\cos \delta$ and three mixing angles
6. If we determine $\cos \delta \neq \pm 1$ \Rightarrow CP is violated!

Direct Analytic Calculation

Disappearance experiments measure various $|U_{\alpha i}|^2$ terms
Suppose 4 are measured: $|U_{e2}|^2, |U_{e3}|^2, |U_{\mu 2}|^2, |U_{\mu 3}|^2$

Actually this gives all 9 magnitudes by unitarity

Direct Analytic Calculation


Disappearance experiments measure various $|U_{\alpha i}|^2$ terms
Suppose 4 are measured: $|U_{e2}|^2, |U_{e3}|^2, |\textcolor{red}{U}_{\mu 2}|^2, |U_{\mu 3}|^2$

Actually this gives all 9 magnitudes by unitarity



$$J_{CP}^2 = |U_{e2}|^2 |U_{\mu 2}|^2 |U_{e3}|^2 |U_{\mu 3}|^2 - \frac{1}{4} (1 - |U_{e2}|^2 - |U_{\mu 2}|^2 - |U_{e3}|^2 - |U_{\mu 3}|^2 + |U_{e2}|^2 |U_{\mu 3}|^2 + |U_{e3}|^2 |U_{\mu 2}|^2)^2$$

Disappearance can tell us if CP is violated, but not if nature prefers ν 's or $\bar{\nu}$'s

Where is $|U_{\mu 2}|^2$?



Where is $|U_{\mu 2}|^2$?



	$\cos \delta$	ROI 1	ROI 2
6.5 yrs ν_μ rates	1	5506	5038
	0	5418	5115
	-1	5334	5193

Approximate size of $|U_{\mu 2}|^2$ signal: (21) sector

- There is no δ information in $|U_{\mu 1}|^2 + |U_{\mu 2}|^2$ (sum of Δ_{31} and Δ_{32} terms)

Approximate size of $|U_{\mu 2}|^2$ signal: (21) sector

- ▶ There is no δ information in $|U_{\mu 1}|^2 + |U_{\mu 2}|^2$ (sum of Δ_{31} and Δ_{32} terms)
- ▶ There is δ information in $|U_{\mu 1}|^2 |U_{\mu 2}|^2$

Approximate size of $|U_{\mu 2}|^2$ signal: (21) sector

- ▶ There is no δ information in $|U_{\mu 1}|^2 + |U_{\mu 2}|^2$ (sum of Δ_{31} and Δ_{32} terms)
- ▶ There is δ information in $|U_{\mu 1}|^2|U_{\mu 2}|^2$
- ▶ This comes from the Δm_{21}^2 term

DUNE and HK can measure Δm_{21}^2 somewhat
PBD, J. Gehrlein [2302.08513](#)

Approximate size of $|U_{\mu 2}|^2$ signal: (21) sector

- ▶ There is no δ information in $|U_{\mu 1}|^2 + |U_{\mu 2}|^2$ (sum of Δ_{31} and Δ_{32} terms)
- ▶ There is δ information in $|U_{\mu 1}|^2|U_{\mu 2}|^2$
- ▶ This comes from the Δm_{21}^2 term

DUNE and HK can measure Δm_{21}^2 somewhat
PBD, J. Gehrlein [2302.08513](#)

- ▶ This term is

$$\approx -4c_{23}^2 (s_{12}^2 c_{12}^2 + s_{23} c_{23} s_{13} \sin 2\theta_{12} \cos 2\theta_{12} \cos \delta) \sin^2 \Delta_{21}$$

$$\approx -2 (0.21 + 0.03 \cos \delta) \left(\frac{\pi}{33} \right)^2$$

$$\Delta m_{21}^2 / |\Delta m_{31}^2| \approx 33$$

Approximate size of $|U_{\mu 2}|^2$ signal: (21) sector

- ▶ There is no δ information in $|U_{\mu 1}|^2 + |U_{\mu 2}|^2$ (sum of Δ_{31} and Δ_{32} terms)
- ▶ There is δ information in $|U_{\mu 1}|^2|U_{\mu 2}|^2$
- ▶ This comes from the Δm_{21}^2 term

DUNE and HK can measure Δm_{21}^2 somewhat
PBD, J. Gehrlein [2302.08513](#)

- ▶ This term is

$$\approx -4c_{23}^2 (s_{12}^2 c_{12}^2 + s_{23} c_{23} s_{13} \sin 2\theta_{12} \cos 2\theta_{12} \cos \delta) \sin^2 \Delta_{21}$$


$$\approx -2 (0.21 + 0.03 \cos \delta) \left(\frac{\pi}{33}\right)^2$$

$$\Delta m_{21}^2 / |\Delta m_{31}^2| \approx 33$$

- ▶ So the probability is large for $\cos \delta = -1$?
- ▶ So the effect is $\sim -0.0005 \cos \delta$?

Approximate size of $|U_{\mu 2}|^2$ signal: (21) sector

- ▶ There is no δ information in $|U_{\mu 1}|^2 + |U_{\mu 2}|^2$
- ▶ There is δ information in $|U_{\mu 1}|^2 |U_{\mu 2}|^2$
- ▶ This comes from the Δm_{21}^2 term



DUNE and HK can measure Δm_{21}^2 somewhat
PBD, J. Gehrlein [2302.08513](#)

- ▶ This term is

$$\approx -4c_{23}^2 (s_{12}^2 c_{12}^2 + s_{23} c_{23} s_{13} \sin 2\theta_{12} \cos 2\theta_{12} \cos \delta) \sin^2 \Delta_{21}$$


$$\approx -2 (0.21 + 0.03 \cos \delta) \left(\frac{\pi}{33}\right)^2$$

$$\Delta m_{21}^2 / |\Delta m_{31}^2| \approx 33$$

- ▶ So the probability is large for $\cos \delta = -1$?
- ▶ So the effect is $\sim -0.0005 \cos \delta$?

Approximate size of $|U_{\mu 2}|^2$ signal: (21) sector

- ▶ There is no δ information in $|U_{\mu 1}|^2 + |U_{\mu 2}|^2$
- ▶ There is δ information in $|U_{\mu 1}|^2 |U_{\mu 2}|^2$
- ▶ This comes from the Δm_{21}^2 term



DUNE and HK can measure Δm_{21}^2 somewhat
PBD, J. Gehrlein [2302.08513](#)

- ▶ This term is

$$\approx -4c_{23}^2 (s_{12}^2 c_{12}^2 + s_{23} c_{23} s_{13} \sin 2\theta_{12} \cos 2\theta_{12} \cos \delta) \sin^2 \Delta_{21}$$

$$\approx -2 (0.21 + 0.03 \cos \delta) \left(\frac{\pi}{33} \right)^2$$

$$\Delta m_{21}^2 / |\Delta m_{31}^2| \approx 33$$

- ▶ So the probability is large for $\cos \delta = -1$?
- ▶ So the effect is $\sim -0.0005 \cos \delta$?

Sign is wrong
Magnitude is ~ 16 too small

Matter effects matter: (21) sector

- ▶ Let's start again at

$$\approx -4c_{23}^2 (s_{12}^2 c_{12}^2 + s_{23} c_{23} s_{13} \sin 2\theta_{12} \cos 2\theta_{12} \cos \delta) \sin^2 \Delta_{21}$$

Matter effects matter: (21) sector

- ▶ Let's start again at

$$\approx -4c_{23}^2 (s_{12}^2 c_{12}^2 + s_{23} c_{23} s_{13} \sin 2\theta_{12} \cos 2\theta_{12} \cos \delta) \sin^2 \Delta_{21}$$

- ▶ Solar splitting modified by

$$\frac{\Delta m_{21}^2 \rightarrow \Delta m_{21}^2 \mathcal{S}_\odot}{\mathcal{S}_\odot \approx \sqrt{(\cos 2\theta_{12} - c_{13}^2 a / \Delta m_{21}^2)^2 + \sin^2 2\theta_{12}} \approx 3.4}$$

at $E = 1.3$ GeV
PBD, S. Parke [1902.07185](#)

Matter effects matter: (21) sector

- ▶ Let's start again at

$$\approx -4c_{23}^2 (s_{12}^2 c_{12}^2 + s_{23} c_{23} s_{13} \sin 2\theta_{12} \cos 2\theta_{12} \cos \delta) \sin^2 \Delta_{21}$$

- ▶ Solar splitting modified by

$$\frac{\Delta m_{21}^2 \rightarrow \Delta m_{21}^2 \mathcal{S}_\odot}{\mathcal{S}_\odot \approx \sqrt{(\cos 2\theta_{12} - c_{13}^2 a / \Delta m_{21}^2)^2 + \sin^2 2\theta_{12}} \approx 3.4}$$

at $E = 1.3$ GeV

PBD, S. Parke [1902.07185](#)

- ▶ Mixing angle is modified

$$\cos 2\theta_{12} = 0.37 \rightarrow \frac{\cos 2\theta_{12} - c_{13}^2 a / \Delta m_{21}^2}{\mathcal{S}_\odot} \approx -0.96 < 0$$

$$a \propto \rho E$$

Matter effects matter: (21) sector

- ▶ Let's start again at

$$\approx -4c_{23}^2 (s_{12}^2 c_{12}^2 + s_{23} c_{23} s_{13} \sin 2\theta_{12} \cos 2\theta_{12} \cos \delta) \sin^2 \Delta_{21}$$

- ▶ Solar splitting modified by

$$\frac{\Delta m_{21}^2 \rightarrow \Delta m_{21}^2 \mathcal{S}_\odot}{\mathcal{S}_\odot \approx \sqrt{(\cos 2\theta_{12} - c_{13}^2 a / \Delta m_{21}^2)^2 + \sin^2 2\theta_{12}} \approx 3.4}$$

at $E = 1.3$ GeV

PBD, S. Parke [1902.07185](#)

- ▶ Mixing angle is modified

$$\cos 2\theta_{12} = 0.37 \rightarrow \frac{\cos 2\theta_{12} - c_{13}^2 a / \Delta m_{21}^2}{\mathcal{S}_\odot} \approx -0.96 < 0$$

- ▶ So the sign is swapped

$$a \propto \rho E$$

$$\sin 2\theta_{12} \cos 2\theta_{12} = 0.37 \rightarrow -0.26$$

Matter effects matter: (21) sector

- ▶ Let's start again at

$$\approx -4c_{23}^2 (s_{12}^2 c_{12}^2 + s_{23} c_{23} s_{13} \sin 2\theta_{12} \cos 2\theta_{12} \cos \delta) \sin^2 \Delta_{21}$$

- ▶ Solar splitting modified by

$$\frac{\Delta m_{21}^2 \rightarrow \Delta m_{21}^2 \mathcal{S}_\odot}{\mathcal{S}_\odot \approx \sqrt{(\cos 2\theta_{12} - c_{13}^2 a / \Delta m_{21}^2)^2 + \sin^2 2\theta_{12}} \approx 3.4}$$

at $E = 1.3$ GeV
PBD, S. Parke [1902.07185](#)

- ▶ Mixing angle is modified

$$\cos 2\theta_{12} = 0.37 \rightarrow \frac{\cos 2\theta_{12} - c_{13}^2 a / \Delta m_{21}^2}{\mathcal{S}_\odot} \approx -0.96 < 0$$

- ▶ So the sign is swapped

$$a \propto \rho E$$

$$\sin 2\theta_{12} \cos 2\theta_{12} = 0.37 \rightarrow -0.26$$

- ▶ Also s_{13} increases in matter $\sim 15\%$: total effect is $0.004 \cos \delta$

Matter effects matter: (21) sector

- ▶ Let's start again at

$$\approx -4c_{23}^2 (s_{12}^2 c_{12}^2 + s_{23} c_{23} s_{13} \sin 2\theta_{12} \cos 2\theta_{12} \cos \delta) \sin^2 \Delta_{21}$$

- ▶ Solar splitting modified by

$$\frac{\Delta m_{21}^2 \rightarrow \Delta m_{21}^2 \mathcal{S}_\odot}{\mathcal{S}_\odot \approx \sqrt{(\cos 2\theta_{12} - c_{13}^2 a / \Delta m_{21}^2)^2 + \sin^2 2\theta_{12}} \approx 3.4}$$

at $E = 1.3$ GeV

PBD, S. Parke [1902.07185](#)

- ▶ Mixing angle is modified

$$\cos 2\theta_{12} = 0.37 \rightarrow \frac{\cos 2\theta_{12} - c_{13}^2 a / \Delta m_{21}^2}{\mathcal{S}_\odot} \approx -0.96 < 0$$

- ▶ So the sign is swapped

$a \propto \rho E$

$$\sin 2\theta_{12} \cos 2\theta_{12} = 0.37 \rightarrow -0.26$$

- ▶ Also s_{13} increases in matter $\sim 15\%$: total effect is $0.004 \cos \delta$
- ▶ This gets us **half** of the effect, and the correct sign

Matter effects matter: (32) sector

- ▶ $\frac{\Delta m_{\mu\mu}^2 L}{4E}$ in matter at the maximum is $\sim \pi$

H. Nunokawa, S. Parke, R. Funchal [hep-ph/0503283](#)
PBD, S. Parke [2401.10326](#)

Matter effects matter: (32) sector

- ▶ $\frac{\Delta m_{\mu\mu}^2 L}{4E}$ in matter at the maximum is $\sim \pi$

H. Nunokawa, S. Parke, R. Funchal [hep-ph/0503283](#)
PBD, S. Parke [2401.10326](#)

- ▶ The Δm_{32}^2 component is a bit off π

Matter effects matter: (32) sector

- ▶ $\frac{\Delta m_{\mu\mu}^2 L}{4E}$ in matter at the maximum is $\sim \pi$

H. Nunokawa, S. Parke, R. Funchal [hep-ph/0503283](#)
PBD, S. Parke [2401.10326](#)

- ▶ The Δm_{32}^2 component is a bit off π
- ▶ Leading order in s_{13} :

$$\begin{aligned} &\approx -4s_{23}^2(c_{12}^2c_{23}^2 - 2s_{13}s_{12}c_{12}s_{23}c_{23}\cos\delta)\sin^2\Delta_{32} \\ &\approx -2 \quad (0.0094 \quad -0.023\cos\delta)0.1 \quad (\text{matter}) \end{aligned}$$

Matter effects matter: (32) sector

- ▶ $\frac{\Delta m_{\mu\mu}^2 L}{4E}$ in matter at the maximum is $\sim \pi$

H. Nunokawa, S. Parke, R. Funchal [hep-ph/0503283](#)
PBD, S. Parke [2401.10326](#)

- ▶ The Δm_{32}^2 component is a bit off π
- ▶ Leading order in s_{13} :

$$\begin{aligned} &\approx -4s_{23}^2(c_{12}^2c_{23}^2 - 2s_{13}s_{12}c_{12}s_{23}c_{23}\cos\delta)\sin^2\Delta_{32} \\ &\approx -2 \quad (0.0094 \quad -0.023\cos\delta)0.1 \quad (\text{matter}) \end{aligned}$$

- ▶ Adds in another $\approx 0.004\cos\delta$ effect

Matter effects matter: (32) sector

- ▶ $\frac{\Delta m_{\mu\mu}^2 L}{4E}$ in matter at the maximum is $\sim \pi$

H. Nunokawa, S. Parke, R. Funchal [hep-ph/0503283](#)
PBD, S. Parke [2401.10326](#)

- ▶ The Δm_{32}^2 component is a bit off π
- ▶ Leading order in s_{13} :

$$\begin{aligned} &\approx -4s_{23}^2(c_{12}^2c_{23}^2 - 2s_{13}s_{12}c_{12}s_{23}c_{23}\cos\delta)\sin^2\Delta_{32} \\ &\approx -2 \quad (0.0094 \quad -0.023\cos\delta)0.1 \quad (\text{matter}) \end{aligned}$$

- ▶ Adds in another $\approx 0.004\cos\delta$ effect
- ▶ Total is $\approx 0.008\cos\delta$ which agrees with exact calculation



Numerical Studies

Inputs are *only*:



- ▶ Daya Bay data for θ_{13} 1809.02261
- ▶ KamLAND data for θ_{12} and Δm_{21}^2 1303.4667
- ▶ JUNO 6 yrs precision sensitivity on θ_{12} , Δm_{21}^2 , Δm_{31}^2 2204.13249
- ▶ DUNE 6.5+6.5 yrs disappearance channels sensitivity only 2103.04797

Also looked at varying JUNO's and DUNE's runtime, and at HK

JUNO and DUNE disappearance Sensitivities



JUNO and HK disappearance Sensitivities



Discussion and Conclusions

- ▶ **Disappearance can discover CPV**
- ▶ Requires two good measurements: JUNO and DUNE/HK
- ▶ Can rule out some values of δ at $> 3\sigma$
- ▶ Analyses already exist but...

Discussion and Conclusions

- ▶ **Disappearance can discover CPV**
- ▶ Requires two good measurements: JUNO and DUNE/HK
- ▶ Can rule out some values of δ at $> 3\sigma$
- ▶ Analyses already exist but...
- ▶ **LBL Experiments should break down δ analyses into app vs. dis**
- ▶ Since systematics are different, provides a good cross check
- ▶ Subject to BSM degeneracies, as are most other oscillation measurements
- ▶ Works in vacuum or matter; matter slightly minimizes HK's effect


Discussion and Conclusions

- ▶ **Disappearance can discover CPV**
- ▶ Requires two good measurements: JUNO and DUNE/HK
- ▶ Can rule out some values of δ at $> 3\sigma$
- ▶ Analyses already exist but...
- ▶ **LBL Experiments should break down δ analyses into app vs. dis**
- ▶ Since systematics are different, provides a good cross check
- ▶ Subject to BSM degeneracies, as are most other oscillation measurements
- ▶ Works in vacuum or matter; matter slightly minimizes HK's effect

HFH!
Happy Beer Bike!

Backups

References



SK [hep-ex/9807003](#)

M. Gonzalez-Garcia, et al. [hep-ph/0009350](#)

M. Maltoni, et al. [hep-ph/0207227](#)

SK [hep-ex/0501064](#)

SK [hep-ex/0604011](#)

T. Schwetz, M. Tortola, J. Valle [0808.2016](#)

M. Gonzalez-Garcia, M. Maltoni, J. Salvado [1001.4524](#)

T2K [1106.2822](#)


D. Forero, M. Tortola, J. Valle [1205.4018](#)

D. Forero, M. Tortola, J. Valle [1405.7540](#)

P. de Salas, et al. [1708.01186](#)

F. Capozzi et al. [2003.08511](#)

Varying Runtime/Power




Improvement requires **both** experiments!

Complex phase in different parameterizations

- ▶ Can relate the complex phase in one parameterization to that in another
- ▶ U_{132} and U_{213} similar to U_{123}
- ▶ δ constrained to $\sim [150^\circ, 210^\circ]$ in U_{231} , U_{312} , U_{321}
- ▶ Bands indicate 3σ uncertainty on θ_{12} , θ_{13} , θ_{23}
- ▶ “50% of possible values of δ ”
⇒ parameterization dependent

DUNE TDR II [2002.03005](#)



Quark mixing

From the PDG, V_{CKM} in the V_{123} parameterization is

$$\theta_{12} = 13.09^\circ \quad \theta_{13} = 0.2068^\circ \quad \theta_{23} = 2.323^\circ \quad \delta_{\text{PDG}} = 68.53^\circ$$



Looks like “large” CPV:

$$\sin \delta_{\text{PDG}} = 0.93 \sim 1$$

yet $J_{\text{CKM}}/J_{\text{max}} = 3 \times 10^{-4}$.

Switch to V_{212} parameterization, $\Rightarrow \delta' = 1^\circ$ and $\sin \delta' = 0.02$.

Standard oscillation parameters



Can see that the combination doesn't like the NO while it does like the IO
IO preferred over NO at $\Delta\chi^2 = 2.3$

CP violation in oscillations

In vacuum at first maximum:

$$P_{\mu e} - \bar{P}_{\mu e} \approx 8\pi J \frac{\Delta m_{21}^2}{\Delta m_{32}^2}$$

$$J \equiv s_{12}c_{12}s_{13}c_{13}^2s_{23}c_{23} \sin \delta$$

C. Jarlskog [PRL 55, 1039 \(1985\)](#)

- ▶ Extracting δ from data requires every other oscillation parameter
- ▶ J requires only Δm_{21}^2 (up to matter effects)


Matter effects are easily accounted for

$$\hat{J} \simeq \frac{J}{\sqrt{(c_{212} - c_{13}^2 a / \Delta m_{21}^2)^2 + s_{212}^2} \sqrt{(c_{213} - a / \Delta m_{ee}^2)^2 + s_{213}^2}}$$

[PBD](#), S. Parke [1902.07185](#)

[PBD](#), H. Minakata, S. Parke [1604.08167](#)


Repeated rotations



	U_{121}	U_{131}	U_{212}	U_{232}	U_{313}	U_{323}
$ U_{e2} $	✓	✓	✓	✓	✗	✗
$ U_{e3} $	✓	✓	✗	✗	✓	✓
$ U_{\mu 3} $	✗	✗	✓	✓	✓	✓

Note that $e^{i\delta}$ must be on first or third rotation


Allowed δ' range



The importance of $\cos \delta$

- ▶ If only $\sin \delta$ is measured \Rightarrow sign degeneracy: $\cos \delta = \pm \sqrt{1 - \sin^2 \delta}$
- ▶ Most flavor models predict $\cos \delta$

J. Gehrlein, et al. [2203.06219](#)



L. Everett, et al. [1912.10139](#)

Rice University: April 2, 2024 39/29